

AD _____

GRANT NUMBER DAMD17-96-1-6242

TITLE: Mechanisms of Murine Mammary Tumorigenesis: Cooperation
Between Tyrosine kinase Receptors and Mutant p53

PRINCIPAL INVESTIGATOR: Archibald Perkins

CONTRACTING ORGANIZATION: Yale University School of Medicine
New Haven, CT 06520-8047

REPORT DATE: August 1998

TYPE OF REPORT: Annual

PREPARED FOR: Commander
U.S. Army Medical Research and Materiel Command
Fort Detrick, Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

19990820 038

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE August 1998	3. REPORT TYPE AND DATES COVERED Annual (1 Aug 97 - 31 Jul 98)
4. TITLE AND SUBTITLE Mechanisms of Murine Mammary Tumorigenesis: Cooperation Between Tyrosine Kinase Receptors and Mutane p53		5. FUNDING NUMBERS DAMD17-96-1-6242
6. AUTHOR(S) Archibald Perkins		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Yale University School of Medicine New Haven, CT 06520-8047		8. PERFORMING ORGANIZATION REPORT NUMBER
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Commander U.S. Army Medical Research and Materiel Command Fort Detrick, Frederick, Maryland 21702-5012		10. SPONSORING/MONITORING AGENCY REPORT NUMBER
11. SUPPLEMENTARY NOTES		19990820 038
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited		12b. DISTRIBUTION CODE

A mouse model for human breast cancer closely mimics the genetic changes that occur in the human disease. Twenty-five to 30% of human breast cancers show amplification and overexpression of the neu (also known as HER2 or ErbB2) gene, and of these, many will have point mutations in p53. The 175H mutation of p53 is the most common p53 mutation in human breast cancers, and is often accompanied by loss of the other allele, arguing that it is not simply acting as a dominant negative. Thus, we have created a useful model for the study of human breast cancer. We are now investigating the mechanism by which this acceleration occurs, taking a variety of approaches that address the role of genomic instability, tumor angiogenesis, and phosphorylation of Neu. Through these approaches we hope to contribute to the current understanding of mammary tumorigenesis as it occurs in the setting of a whole organism.

14. SUBJECT TERMS Breast Cancer			15. NUMBER OF PAGES 25
			16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the U.S. Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

✓ In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Aspaluis
PI - Signature

3/10/99
Date

Annual Report for DAMD17-96-1-6242 for year ending July, 1998.

Table of Contents

Page 1	Front Cover
Page 2	SF 298, Report Documentation Page
Page 3	Foreword
Page 4	Table of Contents
Page 5	Introduction
Page 8	Body
Page 18	Conclusions
Page 18	Figure Legends
Page 19	Literature Cited

Introduction

A central goal of current cancer research is the identification of the genes involved in tumorigenesis, and the definition of the precise role that these genes play in tumor development. Analysis of human breast carcinomas has implicated a number of genes in the genesis of these tumors, including *ErbB2* [43], *HST* and *INT2* [1], *p53* [15], *src* [39], *Rb* [26]. It is suggested by a number of studies that the development of breast cancer in humans requires changes in more than one of these genes, which may in part explain the long latency associated with this disease [15].

ErbB2 encodes a receptor tyrosine kinase related to the receptor for epidermal growth factor (EGFR or ErbB), and is amplified in nearly 30% of human cancers, particularly intraductal carcinomas [17, 44]. Numerous studies suggest that this amplification leads to increased mitogenic signaling in the cell. The importance of this amplification is supported by the finding that 70% of transgenic mice that overexpress rat *ErbB2* in the mammary gland develop mammary carcinomas [11]. However, the latency of tumorigenesis is relatively long (over 200 d), suggesting that other oncogenic events are necessary. Analysis of these tumors revealed small in-frame deletions in the *ErbB2* transgene in 65% of tumors analyzed [42]. These deletions resided in the extracellular domain adjacent to the transmembrane domain, and resulted in activation of *ErbB2* tyrosine kinase activity. These findings indicate that activation of *ErbB2* tyrosine kinase activity plays an important role in the development of these tumors. This is consistent with previous experiments, showing that mice carrying an MMTV-driven rat *ErbB2* transgene with an activating mutation in the transmembrane domain develop multifocal mammary carcinomas with a significantly shorter latency [35].

In 30% of human breast carcinomas, expression of *ERBB2* is associated with the presence of mutant *p53*, suggesting that activated tyrosine kinase receptors cooperate with mutant *p53* in the development of these tumors [15]. *p53* is a multifunctional protein that is involved in the regulation of growth of nearly all cell types within mammalian organisms (reviewed in [23]). The wild type *p53* protein can suppress tumor cell growth [10], and likely functions as a regulatory protein in two capacities: as a key

component of apoptotic pathways within the cell [46]; and as a checkpoint protein to control G1 to S transition in the presence of genotoxic stress [24]. Structural domains of p53 include an amino-terminal transcriptional activation domain, a central DNA binding domain, and a carboxy-terminal domain important for oligomerization (reviewed in [23]). Genetic alterations at the p53 locus are common in human cancers, and are primarily either missense mutations or allele loss [2, 14, 36]. While the majority of human tumors with altered p53 have one allele bearing a missense mutation and one null allele, occasionally tumors are found to have one mutated allele and one normal allele [36]. These findings suggest a progression model in which the initial event is a missense mutation in one p53 allele, leading to a proliferative advantage, and then loss of the other allele, which confers a further selective advantage.

p53 point mutations are highly clustered into four regions that correspond to evolutionarily conserved domains of the protein that function in DNA binding. Some of the most commonly mutated amino acids are those that make direct contact with the DNA [4]. p53 proteins bearing these mutations have been found to have altered DNA binding and transactivation properties [20, 21]. Some mutant proteins fail to activate normal target genes, such as p21, but can activate atypical targets, such as *MDR1* [3]. Thus, certain mutations in p53 may lead to the acquisition of novel and dominant activities within the cell. It is evident from a number of studies that certain missense mutations in p53 function as dominant negative alleles that encode proteins that lack transcriptional activation potential, but retain the ability to oligomerize and thus can pull wildtype p53 into nonfunctional complexes [34]. An example of this is the 135V mutation, which can accelerate tumor development in heterozygous but not nullizygous p53-deficient mice [12]. Other alleles, such as 143A, 175H, 248W, 248Q, 273H, and 281G act as dominant oncogenic alleles, since they can confer new malignant phenotypes upon gene transfer into cells that lack p53 [8, 16]. These phenotypes include the ability to grow in soft agar, and to form invasive tumors in nude mice. The molecular mechanisms that underlie the ability of mutant p53 alleles to induce these changes are unknown.

p53 alterations are common in human breast carcinomas [5, 38]. Missense mutations have been identified at many of the hotspot regions, including codons 175(R to H), and 248(R to Q). 175H represents approximately 8% of all p53 mutations in human breast cancers. These alleles are dominantly oncogenic in cell culture and nude mouse tumorigenicity assays [8, 16]. To obtain a more accurate picture of the effect that the 175H allele has on mammary cell growth, we used transgenic mice in which this allele was targeted to the mammary epithelium using the whey acidic protein (WAP) promoter. It was somewhat surprising to find that, despite high level expression in the mammary gland, mice carrying the WAP-driven p53-175H were not abnormally susceptible to mammary carcinomas - only one mouse developed a mammary carcinoma and this was with a latency of 11 months [27]. These data suggested that this allele is not dominantly oncogenic on its own in this setting, and requires other cooperating events. Indeed, these mice were much more susceptible than nontransgenic control mice to mammary tumors induced by carcinogens that are known to activate Ha-Ras [25, 28, 32]. This suggests that activated Ras is one molecule that can cooperate with p53-172H in this system.

It is known that ErbB2 can initiate a mitogenic signal within the cell, and that this signal utilizes the same pathway as activated ras. This suggested that if p53-172H can cooperate with activated Ras, it may also cooperate with ErbB2. In our previous studies, we demonstrated cooperativity between ErbB2 and p53-172H in the development of mammary carcinomas [29]. These bitransgenic mice constitute a model system that closely mimics the genetic changes in human breast cancers, and that allows for further studies to uncover the mechanism of cooperativity between these two genes.

Body

Experimental Methods

Transgenic Mice The *p53-172H* transgenic mice, in which mutant *p53* transgene was preferentially overexpressed in the mammary epithelium by use of the whey acidic protein (WAP) promoter, were created and characterized as described [27]. *p53* knockout mice were obtained from Tyler Jacks [18]. Unactivated *ErbB2* transgenic mice (line N#202) which contain the wild type rat *ErbB2* gene driven by MMTV have been described previously [11]. All three lines are on an FVB background. *p53/ErbB2* bitransgenic mice were generated by crossing female and male offspring of line 8512 WAP-*p53-172H* transgenic mice to offspring of line N#202 of MMTV-*ErbB2* transgenic mice. Mouse tail DNA from the offspring of this cross was isolated as described previously [29]. The WAP-172H and/or MMTV-*ErbB2* transgenes, and the *p53* knockout allele were identified by PCR as described [18, 29].

Histologic and immunohistologic analysis Mammary glands and mammary tumors were surgically removed, fixed in 10% neutral buffered formalin (ANATECH LTD, Battle Creek, MI) for 6 h, and placed in 70% ethanol until processed. These tissues were embedded in paraffin, and 5 μ m sections were placed on regular slides and stained with hematoxylin and eosin. Immunoperoxidase was performed on deparaffinized tissue sections using standard techniques. To detect vascular endothelial cells, an anti-CD34 antiserum was used (provided by D. Krause, Yale University). To detect phosphorylated neu protein, an antibody directed against a phosphopeptide within the cytoplasmic domain of activated Neu was used (courtesy of M. DiGiovanna).

Spectral Karyotyping. In situ hybridization of fluorescently labeled chromosome-specific probes was performed by Dr. Alan Coleman of the NIH, using techniques described by T. Reid and coworkers [41]

Results and Discussion

1. **Assessing the genetic function of the 172H allele: dominant negative or dominant oncogenic.**

Our previous work describes the creation of a mouse mammary tumor model in which two of the most frequent changes in human breast cancers – amplification of *ErbB2* and a dominant oncogenic mutation of *p53* – have been recapitulated. This model serves to address two important issues in tumor development: the mechanism of cooperation of genes in mammary tumorigenesis, and the effect of dominant oncogenic alleles of *p53* on tumor growth in an in vivo experimental model.

It is known that dominant oncogenic mutants of *p53* such as 175H can cause immortalization of primary cells [40], can cooperate with Ras in transforming primary cells [9, 13], and can enhance the tumorigenic potential of cells lacking *p53* [8]. 175H is particularly potent, being able to induce growth of SAOS-2 cells in agar, where other mutant alleles are not [8]. The rapid kinetics and high efficiency of cooperation in these assays by dominant oncogenic alleles of *p53* indicate a direct effect on tumor cell growth. That these effects can be seen in the absence of endogenous *p53* argues that these alleles are not acting simply as dominant negative alleles, by inactivating wildtype *p53* function. These features of cellular transformation mediated by mutant *p53* alleles suggest that these alleles act not only by interfering with *p53*-dependent functions such as apoptosis, senescence, or genomic instability, [all of which have been suggested as important tumor-promoting sequelae of *p53* loss [19, 31, 33, 45]], but also by exerting a dominant effect on cell growth. The nature of this effect is unknown.

We attempted to determine if the 172H *p53* allele had the capacity to set in a dominant oncogenic manner. We have had difficulty concluding if this is the case. One would predict that a dominant oncogenic allele of *p53* would accelerate tumorigenesis even in the absence of endogenous *p53*. We crossed the *p53*172H allele together with MMTV-Neu onto a *p53*^{-/-} background wishing to compare the latency of mammary tumorigenesis to that seen with: a) 172H and MMTV-neu on a *p53* wt background; b) MMTV-neu on a *p53*^{-/-} background without 172H. Unfortunately, the spectrum of tumors and latency seen in all *p53*^{-/-} mice was the same (lymphomas and sarcomas) regardless of what other transgenes were present.

Table 1.

group	genotype	latency	% MT	%w/sarcoma	%w/lym	#of animals
9	p53-/-	98	0	57	50	14
10	p53-/-+neu	92.5	0	60	60	10
11	p53-/-+172H	99.5	0	75	58	12
12	p53-/-+172H+neu	96.5	0	50	25	4

The results from the p53-/-mice (groups 9,10,11,12) were uninformative concerning the contribution of the 172H transgene to mammary tumorigenesis on a p53 -/- background. We then examined the incidence and latency of mammary tumors on mice heterozygous of endogenous p53 (+/-), bearing either one, both, or neither of the two transgenes (Table 2).

Table 2. Mammary Tumor Incidence and Latency in p53 +/- Mice

(days)		#necropsied	%w/MT*	Latency to MT*
5	p53+/-	4	50%	296
6	p53+/- +neu	10	100%	245
7	p53+/- +172H	13	38%	300
8	p53+/- +neu +172H	9	100%	237

* MT denotes mammary tumor.

The preliminary data from the experiment suggests that in this setting, 172H is not having a significant input, one either the latency or incidence of mammary tumors, with p53+/- genotype, with or without neu.

Table 3. Latency to Mammary Tumor in p53 +/- Mice (in days)

	gp8	gp6
	<u>172H + neu</u>	<u>neu alone</u>
	159	181
	159	221
	215	230
	270	233
median =	229	244 median = 245 d
	246	246
	260	265
	265	304
	277	373
		415

While statistical analysis has not been done, it appears that the latency to tumorigenesis is shorter when the 172H transgene is present. However, it is clear that the affect of the 172H allele on the latency of mammary tumorigenesis is not as dramatic in this setting (p53+/- mice) as it was on the p53+/+ background. One concern here is the level of activity of the WAP promoter driving the 172H allele. The WAP promoter is dependent on the hormonal state of the mouse. It is active from day 10 of gestation through lactation. All these mice were housed with male mice to maintain them in a pregnant and/or lactating status. However, due to a change in personnel, the age at which the mice were placed in mating may have differed from that in our previous experiments (with the p53 +/+ mice). In our previous analysis which compared tumor latency of neu vs. neu+172H on a p53+/+ background, the median age of mammary tumor development was 230 days for neu+ mice, and 150 days for neu + 172H bitransgenic mice. It is not clear that the change in personnel can fully explain the difference in latency between the two experiments, since the age at mating was not consistently later in the more recent experiments with the p53+/- mice. Mice having both the neu+ and 172H transgenes were on the p53+/+ background are currently being maintained in parallel with similar transgenic mice having the p53+/+ genotype (Table 4).

Table 4. Mice currently housed with neu and/or 172H on p53+/+ background

Genotype to MT	# alive	median age	# necropsied	latency
p53+/+ neu+	13	218d	0	-
p53+/+ neu+ 172H	7	197d	1	265
p53+/+ neu+	7	175d	10	245
p53+/+ neu+172H	13	183d	9	237

This more recent cohort of mice (Table 4) should provide pertinent information regarding tumor latency.

To assess the level of 172H transgene expression in these mice, mammary glands and mammary tumors from mice of the various genotypes are being analyzed by Northern blots using a probe for p53. this will provide information about the level of f expression of the 172H transgene, and may provide an explanation for the differences observed between these current experiments and those that we reported on previously (Li et al. 1997).

One curious result is the development of mammary tumors in mice with p53+/- background without the neu transgene. In previous experiments, we found that p53+/+ mice without neu were not significantly susceptible to mammary tumors. This was true for nontransgenic animals as well as 172H transgenic. Our current experiments show that both p53+/- nontransgenic and p53+/- 172H transgenic mice are susceptible to mammary tumors (Table 2). This phenotype is clear with 1721H+,p53+/- mice: of the ten mice that were allowed to age until either lesion development or age>1 year, five developed mammary adenocarcinomas of the MG. One striking finding in these tumors, which had a median latency of 300d, was the presence of abundant fibrovascular stroma surrounding the islands of neoplastic epithelial cells (Figure 1). This histologic feature appears to be specific for this particular genotype, for it has not been seen in any other tumors. This suggests that it is not a nonspecific hormonal effect related to the stage

of these mice in the pregnancy/lactation cycle at which the tumor was harvested. The well-developed stroma seen in the p53+/-, 172H+ tumors is reminiscent of that seen in human breast tumors, a feature that imparts a scirrous quality to breast tumors. The etiology of this intense stromal reaction is unknown. One attractive possibility is that in the absence of a *neu* transgene driving the tumor process, tumorigenesis is instead dependent on mutational activation of other oncogenes. We hypothesize that the 172H transgene is assisting tumor development by increasing the frequency of mutational events and genomic instability. This then leads to activation of cooperating oncogenes, which in this setting, may stimulate stromal proliferation. The identity of such an oncogene(s) is unknown but is likely to encode a growth factor (e.g., FGF, PDGF) or angiogenic growth factor (e.g., VEGF or FGF). One of the five p53+/-, 172H+ tumors was explanted to tissue culture, and can thus be analyzed further for growth factor production and karyotypic abnormalities.

2. Analysis of genomic instability.

In our bitransgenic model, we do not observe the emergence of tumors with kinetics that indicate direct and immediate malignant transformation by coexpression of 172H and *neu*: tumors arise following the second pregnancy rather than the first, and are unifocal, indicating the necessity for other events. This is thus distinct from the cell culture results described above [8], and is likely due to several things, including the lower transforming potential of native *neu* relative to Ras, the presence of endogenous p53 alleles in our transgenic mice, as well as other tumor control mechanisms that exist in the intact animal, such as tumor immunity, the inhibitory influence of surrounding tissue, and the requirement for tumor angiogenesis. Nonetheless, the 172H allele accelerates *neu*-induced tumorigenesis, albeit by an unknown mechanism. We present several possible mechanisms that our bitransgenic model will allow us to address. These models are based on the known or suggested functions of p53, which include an effect on apoptosis, on genome stability, and on transcriptional regulation of cell growth regulatory genes.

The data from cell culture experiments described above suggest a direct effect of 172H on tumor cell growth, and such an effect may indeed play an important role in our system. However, other effects of this allele are also possible. One is that p53-172H increases the likelihood of additional mutational events in genes other than the *neu* transgene in the nonmalignant cells expressing MMTV-*neu*, and thus accelerates tumor formation. One type of genetic alteration known to contribute to mammary tumorigenesis is gene amplification. While an increased frequency of gene amplification is seen in p53 null cells, it is not observed in Li Fraumeni cells (mutated at 184 or 248) that retain one wild-type p53 gene [30]. Since our 172H+*neu* bitransgenic tumors appear by Southern blot analysis to retain (a) wildtype copy or copies of p53 (data not shown), this mechanism may not apply to this model. We are currently assessing the frequency of other types of alterations - e.g., deletions, point mutations - in these bitransgenic tumors.

We are taking three approaches to assessing the contribution of the 172H transgene to mutation frequency and chromosomal instability.

1. Assessing aneuploidy over time in non malignant mammary glands from transgenic and nontransgenic mice.
2. Determination of mutation frequency in mammary tissue of transgenic and nontransgenic mice, using as a mutation sensor a *lacI* indicator gene residing in a lambda phage transgene.
3. Comparative genome hybridization.
4. Spectral karyotyping.

Approach One: We have harvested mammary glands from normal, 172H+, *neu* and 172H+*neu*+mice at various time points, and have processed these for histologic section. H+E slides have been prepared. We will stain these with Feulgen stain, and use image analysis to quantitate the amount of DNA per nucleus.

Approach Two. In collaboration with Dr. Peter Glazer of the Dept. of Therapeutic Radiology at Yale University, we are creating transgenic mice that contain a mutation sensor, which consists of a lambda phage transgene

that contains a lacI gene. Transgenic genomic DNA from these mice can be packaged using standard lambda packaging extracts to yield infectious lambda phage particles. When these are plated on an appropriate indicator strain of E. Coli, one can assess mutation in the lacI gene through blue-white color selection with X-gal. Through this approach, one can determine the mutation frequency in the DNA of any organ within the transgenic mice. We are crossing the lacI transgenic mice with mice containing the 172H and neu alleles to create bi- or tri-transgenic mice that contain lacI. We are now aging these, and will harvest mammary and liver (control) tissues at various time points before and after tumor formation. DNA will be extracted, and will be packaged and assayed for mutation frequency. This experiment will allow us to determine the mutation frequency in nonmalignant and malignant mammary glands in the different genetic backgrounds, which will allow us to assess the influence of the 172H and neu transgenes on mutation frequency.

Approach Three. To document large gains or losses of DNA in tumors arising in mice bearing neu and/or 172H, we are employing comparative genome hybridization (CGH). In this technique, tumor DNA is fluorescently labeled and hybridized to a metaphase spread of normal mouse chromosomes in the presence of normal DNA labeled with a different fluor. Digital photomicroscopy and image analysis are performed to quantitate the relative intensities of the normal and tumor DNA hybridization to the metaphase chromosomes. In this way, regions of DNA overrepresented (amplified) or underrepresented (deleted) in the tumor DNA can be identified. This project is being performed in collaboration with Dr. Alan Coleman at the NIH.

In our lab, we are attempting to perform CGH on nylon filters of mouse BAC libraries instead of on glass slides of metaphase spreads. In this instance, the normal and tumor DNAs are radiolabeled with two different radioactive isotopes (^{32}P and ^{33}P) that can be distinguished on a Phosphoimager. The advantage of this approach is that the regions of increase or loss are present in the tumor can be translated directly into cloned segments of DNA, which can be further analyzed.

Approach Four. In collaboration with Dr. Alan Coleman at NIH we are also analyzing tumors for chromosomal abnormalities by spectral karyotyping (Figure 2). In this experiment, metaphase spreads are prepared from tumor cells, and these are hybridized with a fluorescent probe mix, which contains a probe for each mouse chromosome labeled with a different Fluor. As a result, each mouse chromosome can be distinguished by the emission spectrum of the fluorescence. With this approach, we have documented a chromosomal translocation in a mammary tumor from a mouse containing 172H and neu on a p53+/- background (Figure 2). We are continuing this analysis to determine if the tumors arising in our mice have consistent nonrandom chromosomal abnormalities. Both the number of abnormalities and the identity of the chromosomal region involved will give insight into the degree of genomic instability in mice of different genetic background, and, potentially, the identity of cooperating oncogenic events.

3. Documentation of tumor angiogenesis. An alternative mechanism of p53-172H action in this model is that it may promote other aspects of tumor growth, such as tumor angiogenesis. The finding that mutant, but not wild-type, p53 can synergies with PKC to stimulate vascular endothelial growth factor (VEGF) [22], suggests that the 172H allele could stimulate vascular ingrowth, which is known to be a rate-limiting step in tumorigenesis. To investigate this possibility, we are documenting the density of vessels in mammary tissue and tumors of the different genotypes (no transgene, neu alone, 172H alone, and 172H plus neu). This is being done by immunohistochemical staining for CD34, a cell surface protein present on mouse endothelial cells (Figure 3). This staining is highly specific, and allows for straightforward quantitation of microvessel density in mammary glands and tumors of the different genotypes.

4. Assessment of phosphorylation status of Neu in transgenic tissues, both nonmalignant and malignant. One key feature of neu-induced mammary tumorigenesis is the marked increase in Neu phosphorylation that appears to accompany tumor formation. One potential mechanism by which 172H could accelerate tumor formation is to hasten this increase in phosphorylated Neu. To document the timing of this increase in tumor progression, we are assessing the presence of phosphoNeu at different time

points leading up to tumor formation, using a anti-phosphoneu monoclonal antibody that has been well characterized by Dr. Michael DiGiovanna [6, 7], with whom we are collaborating. We have been able to obtain specific staining with low background (Figure 3), and are now embarking on the analysis of tissue sections from a panel of nonmalignant and malignant mammary tissue specimens obtained at different ages in our transgenic mice.

5. Identification of cooperating oncogenes by MMTV proviral tagging. To further understand the latency of tumorigenesis in our transgenic mice, we have initiated an experiment to identify cooperating oncogenes by MMTV proviral tagging. Two separate experiments are being carried out. In the first, (Table 5), the 172H transgene was put onto a C3H background by backcrossing, and then a transgenic male was crossed to a C3H MMTV+ female mouse, which presumably passed MMTV onto the pups through the milk (this has yet to be confirmed). We have a cohort of 34 mice with both MMTV and 172H, and 13 mice with MMTV alone.

Table 5.

Genotype	MMTV + 172H	MMTV alone
Number in Cohort	34	13

In a second cross, we are creating 172H/neu bitransgenic mice with MMTV by mating bitransgenic male FVB mice with C3H MMTV+ female mice. This cross generates FVB x C3H F1 hybrid mice, with either one, neither, or both transgenes. To date we have the following numbers in each cohort:

Table 6.

Genotype	Neu -, 172H -	neu -, 172H +	neu +, 172H -	neu +, 172H +
Number in Cohort	4	3	4	8

As control cohorts, we are also establishing F1 hybrid mice without MMTV, both transgenic and nontransgenic. To date, none of the mice in the MMTV proviral tagging has developed a mammary tumor. We will continue to monitor these mice.

Conclusions. We have created a mouse model for human breast cancer closely mimics the genetic changes that occur in the human disease. Twenty-five to 30% of human breast cancers show amplification and overexpression of *ErbB2* gene, and of these, many will have point mutations in p53 [15]. The 175H mutation is the most common p53 mutation in human breast cancers, and is often accompanied by loss of the other allele, arguing that it is not simply acting as a dominant negative [37]. Thus, we have created a useful model for the study of human breast cancer. We are now investigating the mechanism by which this acceleration occurs, taking a variety of approaches that address the role of genomic instability, tumor angiogenesis, and phosphorylation of Neu. Through these approaches we hope to contribute to the current understanding of mammary tumorigenesis as it occurs in the setting of a whole organism.

References

1. Ali, I.U., G. Merlo, R. Gallahan, and R. Lidereau. 1989. The amplification unit on chromosome 11q13 in aggressive primary human breast tumors entails the bcl-1, int-2 and hst loci. *Oncogene*. 4:89-92.
2. Caron de Fromental, C. and T. Soussi. 1992. TP53 tumor suppressor gene: a model for investigating human mutagenesis. *Genes Chrom Cancer*. 4:1-15.
3. Chin, K.-V., K. Ueda, I. Pastan, and M.M. Gottesman. 1992. Modulation of activity of the promoter of the human MDR1 gene by Ras and p53. *Science*. 255:459-462.
4. Cho, Y., S. Gorina, P.D. Jeffrey, and N.P. Pavletich. 1994. Crystal structure of a p53 tumor suppressor-DNA complex: understanding tumorigenic mutations. *Science*. 265:346-355.
5. Davidoff, A.M., P.A. Humphrey, J.D. Iglehart, and J.R. Marks. 1991. Genetic basis for p53 overexpression in human breast cancer. *Proc Natl Acad Sci USA*. 88:5006-5010.
6. DiGiovanna, M., M. Lerman, R. Coffey, W. Muller, R. Cardiff, and D. Stern. 1998. Active Signaling by Neu in transgenic mice. *Oncogene*. 17:1877-1884.
7. DiGiovanna, M. and D. Stern. 1995. Activation state specific monoclonal antibody detects tyrosine phosphorylated p185neu/erbB-2 in a subset of human breast tumors overexpressing this receptor. *Cancer Res*. 55:1946-1955.
8. Dittmer, D., S. Pati, G. Zambetti, S. Chu, A.K. Teresky, M. Moore, C. Finlay, and A.J. Levine. 1993. Gain of function mutations in p53. *Nature Genetics*. 4:42-46.
9. Eliyahu, D., A. Raz, P. Gruss, D. Givol, and M. Oren. 1984. Participation of p53 cellular tumor antigen in transformation of normal embryonic cells. *Nature*. 312:646-649.
10. Finlay, C.A., P.W. Hinds, and A.J. Levine. 1989. The p53 proto-oncogene can act as a suppressor of transformation. *Cell*. 57:1083-1093.
11. Guy, S.T., M.A. Webster, M. Schaller, T.J. Parsons, R.D. Cardiff, and W.J. Muller. 1992. Expression of the neu protooncogene in the mammary epithelium of transgenic mice induces metastatic disease. *Proc Natl Acad Sci, USA*. 89:10578-10582.
12. Harvey, M., H. Vogel, D. Morris, A. Bradley, A. Bernstein, and L.A. Donehower. 1995. A mutant p53 transgene accelerates tumour development in heterozygous but not nullizygous p53-deficient mice. *Nature Genetics*. 9:305-311.
13. Hinds, P.W., C.A. Finlay, R.S. Quartin, S.J. Baker, E.R. Fearon, B. Vogelstein, and A.J. Levine. 1990. Mutant p53 DNA clones from human colon carcinomas cooperate with ras in transforming primary rat cells: a comparison of the "hot spot" mutant phenotypes. *Cell Growth Differ*. 1:571-580.
14. Hollstein, M., D. Sidransky, B. Vogelstein, and C.C. Harris. 1991. p53 mutations in human cancers. *Science*. 253:49-53.
15. Horak, E., K. Smith, L. Bromley, S. LeJeune, M. Greenall, D. Lane, and A.L. Harris. 1991. Mutant p53, EGF receptor and c-erbB-2 expression in human breast cancer. *Oncogene*. 6:2277-2284.
16. Hsiao, M., J. Low, E. Dorn, D. Ku, P. Pattengale, J. Yeargin, and M. Haas. 1994. Gain-of-function mutations of the p53 gene induce

- lymphohematopoietic mteastatic potential and tissue invasiveness. *Am J Pathol.* 145:702-714.
17. Hynes, N.E. and D.F. Stern. 1994. The biology of *erbB-2/neu/HER-2* and its role in cancer. *Biochemica et Biophysica Acta Reviews on Cancer.* 1198:165-184.
 18. Jacks, T., L. Remington, B. Williams, E. Schmitt, S. Halachmi, R. Bronson, and R. Weinberg. 1994. Tumor spectrum analysis in p53-mutant mice. *Current Biology.* 4:1-7.
 19. Kastan, M.B., O. Onyekwere, D. Sidransky, B. Vogelstein, and R. Craig. 1991. Participation of p53 protein in the cellular response to DNA damage. *Cancer Res.* 51:6304-6311.
 20. Kern, S.E., K. K.W., B. S.J., N. J.M., R. V., L. A.J., F. P., P. C., and V. B. 1991. Mutated p53 binds DNA abnormally in vitro. *Oncogene.* 6:131-136.
 21. Kern, S.E., J.A. Pietenpol, S. Thiagalingam, A. Seymour, K.W. Kinzler, and B. Vogelstein. 1992. Oncogenic forms of p53 inhibit p53-regulated gene expression. *Science.* 256:827-832.
 22. Kieser, A., H.A. Weich, G. Brandner, D. Marme, and W. Kolch. 1994. Mutant p53 potentiates protein kinase C induction of vascular endothelial growth factor expression. *Oncogene.* 9:963-969.
 23. Ko, L.J. and C. Prives. 1996. p53: puzzle and paradigm. *Genes Dev.* 10:1054-1072.
 24. Kuerbitz, S.J., B.S. Plunkett, W.V. Walsh, and M.B. Kastan. 1992. Wild-type p53 is a cell cycle checkpoint determinant following irradiation. *Proc Natl Acad Sci USA.* 89:7491-7495.
 25. Kumar, R., Medina, D. & Sukumar, S. 1990. Activation of H-Ras oncogene in preneoplastic mouse mammary tissues. *Oncogene.* 5:1271-1277.
 26. Lee, E.Y.-H., H. To, J.Y. Shew, R. Bookstein, P. Scully, and W.-H. Lee. 1988. Inactivation of the retinoblastoma susceptibility gene in human breast cancers. *Science.* 241:218-221.
 27. Li, B., D. Medina, and J.M. Rosen. 1996. Acceleration of DMBA-induced mammary tumorigenesis with a 172 arg to his mutant p53. in preparation.
 28. Li, B., K. Murphy, R. Laucirica, F. Kittrell, D. Medina, and J. Rosen. 1998. A transgenic mouse model for mammary carcinogenesis. *Oncogene.* 16:997-1007.
 29. Li, B., J. Rosen, J. McMenamin-Balano, W. Muller, and A. Perkins. 1997. *Neu/ErbB2* cooperates with p53 -72H during murine mammary tumorigenesis in transgenic mice. *Mol Cell Biol.* 17:3155-3163.
 30. Livingstone, L.R., A. White, J. Sprouse, E. Livanos, T. Jacks, and T.D. Tlsty. 1992. Altered cell cycle arrest and gene amplification potential accompany loss of wild-type p53. *Cell.* 70:923-935.
 31. Lowe, S.W., E.M. Schmitt, S.W. Smith, B.A. Osborne, and T. Jacks. 1993. p53 is required for radiation-induced apoptosis in mouse thymocytes. *Nature.* 362:847-849.
 32. Medina, D. 1974. Mammary tumorigenesis in chemical carcinogen-treated mice. I. Incidence in Balb-c and C57BL mice. *J. Natl. Cancer Institute.* 53:213-221.
 33. Metz, T., A.W. Harris, and J.M. Adams. 1995. Absence of p53 allows direct immortalization of hematopoietic cells by the *myc* and *raf* oncogenes. *Cell.* 82:29-36.
 34. Milner, J. and E.A. Medcalf. 1991. Cotranslation of activated mutant p53 with wild type drives the wild-type p53 protein into the mutant conformation. *Cell.* 65:765-774.

35. Muller, W.J., E. Sinn, P.K. Pattengale, R. Wallace, and P. Leder. 1988. Single-step induction of mammary adenocarcinoma in transgenic mice bearing the activated c-neu oncogene. *Cell*. 54:105-115.
36. Nigro, J.M., S.J. Baker, A.C. Preisinger, J.M. Jessup, R. Hostetter, K. Cleary, S.H. Bigner, N. Davidson, S. Baylin, P. Devilee, T. Glover, F.S. Collins, A. Weston, R. Modali, C.C. Harris, and B. Vogelstein. 1989. Mutations in the p53 gene occur in diverse human tumor types. *Nature*. 342:705-708.
37. Ozbun, M.A. and J.S. Butel. 1995. Tumor suppressor p53 mutations and breast cancer: a critical analysis. *Advances in Cancer Research*. 66:71-141.
38. Prosser, J., A.M. Thompson, G. Cranson, and H.J. Evans. 1990. Evidence that p53 behaves as a tumour suppressor gene in sporadic breast tumours. *Oncogene*. 5:1573-1579.
39. Rosen, N., J.B. Bolen, A.M. Schwartz, P. Cohen, V. DeSeau, and M.A. Israel. 1986. Analysis of pp60 c-src protein kinase activity in human tumor cell lines and tissues. *J. Biol. Chem*. 261:13754-13759.
40. Rovinski, B. and S. Benchimol. 1988. immortalization of rat embryo fibroblasts by the cellular p53 oncogene. *Oncogene*. 2:445-452.
41. Schrock, E., S. du Manoir, T. Veldman, B. Schoell, J. Wienberg, M. Ferguson-Smith, Y. Ning, D. Ledbetter, I. Bar-Am, D. Soenksen, Y. Garini, and T. Reid. 1996. Multicolor spectral karyotyping of human chromosomes. *Science*. 273:494-497.
42. Siegel, P.M., D.L. Dankort, W.R. Hardy, and W.J. Muller. 1994. Novel activating mutations in the neu proto-oncogene involved in induction of mammary tumors. *Mol Cell Biol*. 14:7068-7077.
43. Slamon, D.J., G.M. Clark, S.G. Wong, W.J. Levin, A. Ullrich, and R.L. McGuire. 1987. Human breast cancer: correlation of relapse and survival with amplification of the HER-2/Neu oncogene. *Science*. 235:177-182.
44. Slamon, D.J., W. Godolphin, L.A. Jones, J.A. Holt, S.C. Wong, D.E. Keith, W.J. Levin, S.G. Stuart, J. Udore, A. Ullrich, and M.F. Press. 1989. Studies of the HER-2/neu protooncogene in human breast and ovarian cancer. *Science*. 244:707-712.
45. Symonds, H., L. Krall, L. Remington, M. Saenz-Robles, S. Lowe, T. Jacks, and T. Van Dyke. 1994. p53-dependent apoptosis suppresses tumor growth and progression in vivo. *Cell*. 78:703-711.
46. Yonish-Rouach, E., D. Resnitsky, J. Lotem, L. Sachs, A. Kimchi, and M. Oren. 1991. Wildtype p53 induces apoptosis of myeloid leukemic cells that is inhibited by interleukin-6. *Nature*. 352:345-347.

Figure Legends.

Fig. 1. Histology of tumors arising in the mammary glands of mice with various genotypes. Shown are photomicrographs of H+E stained sections of mammary tumors arising in mice with the genotypes indicated to the top of each panel.


Fig. 2. Spectral karyotyping of tumor #5132 (p53+/-; neu+; 172H+). Left panel: histopathology (H+E stain); right panel: spectral karyotyping. Arrow denotes translocation involving chromosomes 5 and 7.

Fig. 3. Immunoperoxidase analysis of mammary glands (top panels) and mammary tumors (bottom four panels) performed with the antibodies indicated. Ctl denotes control immunoperoxidase reaction with no primary antibody.


Gr 4 #5545 p53+/+;neu+;p53172H+




Gr 5 #5035 p53+/-;neu-;p53172H-



Gr 7 #4923 p53+/-;neu-;p53172H+



Gr 7 #5033 p53+/-;neu-;p53172H+



Gr 6 #5066 p53+/-;neu+;p53172H-



Gr 8 #5132 p53+/-;neu+;p53172H+






Figure 1




p53+/-; neu+; 172H+


Figure 2




CD34 ImPox



Ctl ImPox




P-Neu ImPox



Ctl ImPox



P-Neu ImPox



Ctl ImPox

Figure 3